arXiv:1207.7114v1 [math.CA] 30 Jul 2012

NEW ESTIMATES ON GENERALIZATION OF SOME INTEGRAL INEQUALITIES FOR $s-$CONVEX FUNCTIONS AND THEIR APPLICATIONS

İMDAT İŞCAN

Abstract. In this paper, a new identity for differentiable functions is derived. Thus we can obtain new estimates on generalization of Hadamard,Ostrowski and Simpson type inequalities for functions whose derivatives in absolute value at certain power are $s-$convex (in the second sense). Some applications to special means of real numbers are also given.

1. Introduction

Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a convex function defined on the interval I of real numbers and $a, b \in I$ with $a < b$. The following inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a}\int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2}. \tag{1.1}$$

holds. This double inequality is known in the literature as Hermite-Hadamard integral inequality for convex functions [6]. Note that some of the classical inequalities for means can be derived from (1.1) for appropriate particular selections of the mapping f. Both inequalities hold in the reversed direction if f is concave.

Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a mapping differentiable in I°, the interiorof I, and let $a, b \in I^{\circ}$ with $a < b$. If $|f'(x)| \leq M$, $x \in [a,b]$, then we the following inequality holds

$$\left| f(x) - \frac{1}{b-a}\int_a^b f(t)dt \right| \leq \frac{M}{b-a}\left[\frac{(x-a)^2+(b-x)^2}{2}\right]$$

for all $x \in [a,b]$. The constant $\frac{1}{4}$ is the best possible in the sense that it cannot be replaced by a smaller one. This result is known in the literature as the Ostrowski inequality [7].

The following inequality is well known in the literature as Simpson's inequality .

Date: July 21, 2012.

2000 *Mathematics Subject Classification.* 26A51, 26D15.

Key words and phrases. Hermite-Hadamard inequality, Simpson type inequality, Ostrowski type inequality, trapezoid type inequality, midpoint type inequality, $s-$convex function, convex function.

Let $f:[a,b]\rightarrow\mathbb{R}$ be a four times continuously differentiable mapping on (a,b) and $\left\|f^{(4)}\right\|_{\infty}=\sup\limits_{x\in(a,b)}\left|f^{(4)}(x)\right|<\infty$. Then the following inequality holds:

$$\left|\frac{1}{3}\left[\frac{f(a)+f(b)}{2}+2f\left(\frac{a+b}{2}\right)\right]-\frac{1}{b-a}\int\limits_{a}^{b}f(x)dx\right|\leq\frac{1}{2880}\left\|f^{(4)}\right\|_{\infty}(b-a)^{2}.$$

In recent years many authors have studied error estimations for Simpson's inequality; for refinements, counterparts, generalizations and new Simpson's type inequalities, see [1, 11, 12, 13] and therein.

In [4], Breckner introduced s-convex functions as a generalization of convex functions as follows:

Definition 1. *Let $s\in(0,1]$ be a fixed real number. A function $f:[0,\infty)\rightarrow[0,\infty)$ is said to be $s-$convex (in the second sense),or that f belongs to the class K_{s}^{2}, if*

$$f\left(tx+(1-t)y\right)\leq t^{s}f(x)+(1-t)^{s}f(y) \tag{1.2}$$

for all $x,y\in[0,\infty)$ and $t\in[0,1]$.

If inequality (1.2) is reversed, then f is said to be $s-$concave (in the second sense). Of course, s-convexity means just convexity when s = 1.

In [5], Dragomir and Fitzpatrick proved a variant of Hadamard's inequality which holds for $s-$convex functions in the second sense.

Theorem 1. *Suppose that $f:[0,\infty)\rightarrow[0,\infty)$ is an $s-$convex function in the second sense, where $s\in(0,1)$, and let $a,b\in[0,\infty)$, $a<b$.If $f\in L[a,b]$ then the following inequalities hold*

$$2^{s-1}f\left(\frac{a+b}{2}\right)\leq\frac{1}{b-a}\int\limits_{a}^{b}f(x)dx\leq\frac{f(a)+f(b)}{s+1}. \tag{1.3}$$

Both inequalities hold in the reversed direction if f is $s-$concave. The constant $k=\frac{1}{s+1}$ is the best possible in the second inequality in (1.3).

2. Main results

In order to prove our main theorems, we need the following Lemma.

Lemma 1. *Let $f:I\subseteq\mathbb{R}\rightarrow\mathbb{R}$ be a differentiable mapping on I° such that $f'\in L[a,b]$, where $a,b\in I$ with $a<b$ and $\theta,\lambda\in[0,1]$. Then the following equality holds:*

$$\begin{aligned}&(1-\theta)\left(\lambda f(a)+(1-\lambda)f(b)\right)+\theta f((1-\lambda)a+\lambda b)-\frac{1}{b-a}\int\limits_{a}^{b}f(x)dx\\ =\;&(b-a)\left[-\lambda^{2}\int\limits_{0}^{1}(t-\theta)f'\left(ta+(1-t)\left[(1-\lambda)a+\lambda b\right]\right)dt\right.\\ &\left.+(1-\lambda)^{2}\int\limits_{0}^{1}(t-\theta)f'\left(tb+(1-t)\left[(1-\lambda)a+\lambda b\right]\right)dt\right].\end{aligned}$$

Proof. Firstly suppose that $\lambda \in (0,1)$ and let $C = (1-\lambda)\,a + \lambda b$.

$$I = -\lambda^2 \int_0^1 (t-\theta)\, f'\left(ta + (1-t)\,C\right) dt + (1-\lambda)^2 \int_0^1 (t-\theta)\, f'\left(tb + (1-t)\,C\right) dt$$

integrating by parts, we get

$$\begin{aligned} I &= (t-\theta)\,\frac{\lambda f\left(ta + C\,(1-t)\right)}{b-a}\Bigg|_0^1 - \frac{\lambda}{b-a}\int_0^1 f\left(ta + C\,(1-t)\right) dt \\ &\quad + (t-\theta)\,\frac{(1-\lambda)\, f\left(tb + C\,(1-t)\right)}{b-a}\Bigg|_0^1 - \frac{(1-\lambda)}{b-a}\int_0^1 f\left(tb + C\,(1-t)\right) dt \\ &= \lambda\,(1-\theta)\,\frac{f\,(a)}{b-a} + \lambda\theta\frac{f\,(C)}{b-a} + (1-\lambda)\,(1-\theta)\,\frac{f\,(b)}{b-a} + (1-\lambda)\,\theta\frac{f\,(C)}{b-a} \\ &\quad - \frac{1}{b-a}\left[\lambda\int_0^1 f\left(ta + C\,(1-t)\right) dt + (1-\lambda)\int_0^1 f\left(tb + C\,(1-t)\right) dt\right] \end{aligned}$$

Setting $x = ta + C\,(1-t)\,,\ dx = \lambda\,(a-b)\,dt$,and $x = tb + C\,(1-t)\,,\ dx = \mu\,(b-a)\,dt$ respectively, we obtain

$$\begin{aligned} &\lambda\int_0^1 f\left(ta + C\,(1-t)\right) dt + (1-\lambda)\int_0^1 f\left(tb + C\,(1-t)\right) dt \\ &= \frac{1}{b-a}\left[\int_a^C f\,(x)\,dx + \int_C^b f\,(x)\,dx\right] = \frac{1}{b-a}\int_a^b f\,(x)\,dx \end{aligned}$$

and so we have

$$(b-a)\, I = (1-\theta)\left(\lambda f(a) + (1-\lambda)\, f(b)\right) + \theta f(C) - \frac{1}{b-a}\int_a^b f(x)dx$$

which gives the desired representation (2.1).

Secondly suppose that $\lambda \in \{0,1\}$. The identities

$$(1-\theta)\, f(b) + \theta f(a) - \frac{1}{b-a}\int_a^b f(x)dx = (b-a)\int_0^1 (t-\theta)\, f'\left(tb + (1-t)\,a\right) dt$$

and

$$(1-\theta)\, f(a) + \theta f(b) - \frac{1}{b-a}\int_a^b f(x)dx = (b-a)\int_0^1 (\theta-t)\, f'\left(ta + (1-t)\,b\right) dt.$$

can be proved by performing an integration by parts in the integrals from the right side and changing the variable. $\square$

Theorem 2. *Let $f : I \subseteq [0,\infty) \to \mathbb{R}$ be a differentiable mapping on I° such that $f' \in L[a,b]$, where $a,b \in I^{\circ}$ with $a < b$ and $\theta, \lambda \in [0,1]$. If $|f'|^q$ is $s-$convex on $[a,b]$, $q \geq 1$, then the following inequality holds:*

$$\left|(1-\theta)\left(\lambda f(a)+(1-\lambda)f(b)\right)+\theta f((1-\lambda)a+\lambda b)-\frac{1}{b-a}\int_a^b f(x)dx\right|$$

$$\begin{aligned}
&\leq (b-a)\, A_1^{1-\frac{1}{q}}(\theta)\left\{\lambda^2\left[\left|f'(a)\right|^q A_2(\theta,s)+\left|f'(C)\right|^q A_3(\theta,s)\right]^{\frac{1}{q}}\right. \\
&\qquad \left. +(1-\lambda)^2\left[\left|f'(b)\right|^q A_2(\theta,s)+\left|f'(C)\right|^q A_3(\theta,s)\right]^{\frac{1}{q}}\right\} \qquad (2.1)
\end{aligned}$$

where

$$\begin{aligned}
A_1(\theta) &= \theta^2-\theta+\frac{1}{2} \\
A_2(\theta,s) &= \frac{2\theta^{s+2}}{(s+1)(s+2)}-\frac{\theta}{s+1}+\frac{1}{s+2}, \\
A_3(\theta,s) &= \frac{2(1-\theta)^{s+2}}{(s+1)(s+2)}-\frac{1-\theta}{s+1}+\frac{1}{s+2}.
\end{aligned}$$

and $C=(1-\lambda)a+\lambda b$.

Proof. Suppose that $q \geq 1$ and $C=(1-\lambda)a+\lambda b$. From Lemma 1 and using the well known power mean inequality, we have

$$\left|(1-\theta)\left(\lambda f(a)+(1-\lambda)f(b)\right)+\theta f(C)-\frac{1}{b-a}\int_a^b f(x)dx\right| \leq (b-a)$$

$$\left[\lambda^2\int_0^1 |t-\theta|\left|f'(ta+(1-t)C)\right|dt+(1-\lambda)^2\int_0^1 |t-\theta|\left|f'(tb+(1-t)C)\right|dt\right]$$

$$\begin{aligned}
&\leq (b-a)\left\{\lambda^2\left(\int_0^1|t-\theta|dt\right)^{1-\frac{1}{q}}\left(\int_0^1|t-\theta|\left|f'(ta+(1-t)C)\right|^q dt\right)^{\frac{1}{q}}\right. \\
&\qquad \left.+(1-\lambda)^2\left(\int_0^1|t-\theta|dt\right)^{1-\frac{1}{q}}\left(\int_0^1|t-\theta|\left|f'(tb+(1-t)C)\right|^q dt\right)^{\frac{1}{q}}\right\}. \qquad (2.2)
\end{aligned}$$

Since $|f'|^q$ is $s-$convex on $[a,b]$, we know that for $t \in [0,1]$

$$\left|f'(ta+C(1-t))\right|^q \leq t^s\left|f'(a)\right|^q+(1-t)^s\left|f'(C)\right|^q \qquad (2.3)$$

and

$$\left|f'(tb+C(1-t))\right|^q \leq t^s\left|f'(b)\right|^q+(1-t)^s\left|f'(C)\right|^q. \qquad (2.4)$$

Hence, by simple computation

$$\begin{aligned}
&\int_0^1 |t-\theta| \left|f'\left(ta+(1-t)C\right)\right|^q dt \\
\leq\;& \int_0^1 |t-\theta| \left[t^s \left|f'(a)\right|^q + (1-t)^s \left|f'(C)\right|^q\right] dt \\
=\;& \left|f'(a)\right|^q \int_0^1 |t-\theta| t^s dt + \left|f'(C)\right|^q \int_0^1 |t-\theta| (1-t)^s dt \\
=\;& \left|f'(a)\right|^q A_2(\theta,s) + \left|f'(C)\right|^q A_3(\theta,s) \qquad (2.5)
\end{aligned}$$

$$\begin{aligned}
&\int_0^1 |t-\theta| \left|f'\left(tb+(1-t)C\right)\right|^q dt \\
\leq\;& \int_0^1 |t-\theta| \left[t^s \left|f'(b)\right|^q + (1-t)^s \left|f'(C)\right|^q\right] dt \\
=\;& \left|f'(b)\right|^q \int_0^1 |t-\theta| t^s dt + \left|f'(C)\right|^q \int_0^1 |t-\theta| (1-t)^s dt \\
=\;& \left|f'(b)\right|^q A_2(\theta,s) + \left|f'(C)\right|^q A_3(\theta,s) \qquad (2.6)
\end{aligned}$$

and

$$\int_0^1 |t-\theta| \, dt = \theta^2 - \theta + \frac{1}{2}. \qquad (2.7)$$

Thus, using (2.5),(2.6) and (2.7) in (2.2), we obtain the inequality (2.1). This completes the proof. □

Corollary 1. *Under the assumptions of Theorem 2 with* $q=1$, *we have*

$$\begin{aligned}
&\left|(1-\theta)\left(\lambda f(a)+(1-\lambda)f(b)\right)+\theta f\left((1-\lambda)a+\lambda b\right)-\frac{1}{b-a}\int_a^b f(x)dx\right| \\
\leq\;& (b-a)\left\{A_2(\theta,s)\left(\lambda^2\left|f'(a)\right|+(1-\lambda)^2\left|f'(b)\right|\right)+A_3(\theta,s)\left(2\lambda^2-2\lambda+1\right)\left|f'(C)\right|\right\}
\end{aligned}$$

where $A_2(\theta,s)$ *and* $A_3(\theta,s)$ *are defined as in Theorem 2.*

Corollary 2. *Under the assumptions of Theorem 2 with* $s=1$ *we have*

$$\begin{aligned}
&\left|(1-\theta)\left(\lambda f(a)+(1-\lambda)f(b)\right)+\theta f(C)-\frac{1}{b-a}\int_a^b f(x)dx\right| \\
\leq\;& (b-a)\,A_1^{1-\frac{1}{q}}(\theta)\left\{\lambda^2\left[\left|f'(a)\right|^q A_2(\theta,1)+\left|f'(C)\right|^q A_3(\theta,1)\right]^{\frac{1}{q}}\right. \\
&\left.+(1-\lambda)^2\left[\left|f'(b)\right|^q A_2(\theta,1)+\left|f'(C)\right|^q A_3(\theta,1)\right]^{\frac{1}{q}}\right\}
\end{aligned}$$

where

$$\begin{aligned} A_1(\theta) &= \theta^2-\theta+\frac{1}{2} \\ A_2(\theta,1) &= \frac{1+\theta^3}{3}-\frac{\theta}{2}, \\ A_3(\theta,1) &= \frac{1+(1-\theta)^3}{3}-\frac{1-\theta}{2}, \end{aligned}$$

and $C=(1-\lambda)a+\lambda b$.

Corollary 3. *Under the assumptions of Theorem 2 with* $\theta=1$, *then we have following generalized midpoint type inequality*

$$\begin{aligned} &\left| f((1-\lambda)a+\lambda b)-\frac{1}{b-a}\int_a^b f(x)dx \right| \\ \leq\ & \frac{b-a}{2}\left(\frac{2}{(s+1)(s+2)}\right)^{\frac{1}{q}}\left\{\lambda^2\left(|f'(a)|^q+(s+1)|f'(C)|^q\right)^{\frac{1}{q}}\right. \\ & \left. +(1-\lambda)^2\left(|f'(b)|^q+(s+1)|f'(C)|^q\right)^{\frac{1}{q}}\right\}, \end{aligned}$$

where $C=(1-\lambda)a+\lambda b$.

Corollary 4. *Under the assumptions of Theorem 2 with* $\theta=1$, *if* $|f'(x)|\leq M$, $x\in[a,b]$, *then we have the following Ostrowski type inequality*

$$\left| f(x)-\frac{1}{b-a}\int_a^b f(u)du \right| \leq M\left(\frac{2}{s+1}\right)^{\frac{1}{q}}\left[\frac{(x-a)^2+(b-x)^2}{2(b-a)}\right] \tag{2.8}$$

for each $x\in[a,b]$.

Proof. For each $x\in[a,b]$, there exist $\lambda_x\in[0,1]$ such that $x=(1-\lambda_x)a+\lambda_x b$. Hence we have $\lambda_x=\frac{x-a}{b-a}$ and $1-\lambda_x=\frac{b-x}{b-a}$. Therefore for each $x\in[a,b]$, from the inequality (2.1) we obtain the inequality (2.8). □

Remark 1. *We note that the inequality (2.8) is the same of the inequality in* [2, Theorem 4].

Corollary 5. *Under the assumptions of Theorem 2 with* $\theta=0$, *then we have following generalized trapezoid type inequality*

$$\begin{aligned} &\left| \lambda f(a)+(1-\lambda)f(b)-\frac{1}{b-a}\int_a^b f(x)dx \right| \\ \leq\ & \frac{b-a}{2}\left(\frac{2}{(s+1)(s+2)}\right)^{\frac{1}{q}}\left\{\lambda^2\left((s+1)|f'(a)|^q+|f'(C)|^q\right)^{\frac{1}{q}}\right. \\ & \left. +(1-\lambda)^2\left((s+1)|f'(b)|^q+|f'(C)|^q\right)^{\frac{1}{q}}\right\}, \end{aligned}$$

where $C=(1-\lambda)a+\lambda b$.

Corollary 6. *Under the assumptions of Theorem 2 with* $\lambda = \frac{1}{2}$ *and* $\theta = \frac{2}{3}$*, then we have the following Simpson type inequality*

$$\begin{aligned}
&\left|\frac{1}{6}\left[f(a)+4f\left(\frac{a+b}{2}\right)+f(b)\right]-\frac{1}{b-a}\int_a^b f(x)dx\right| \\
\leq\ & \frac{b-a}{4}\left(\frac{5}{18}\right)^{1-\frac{1}{q}}\left\{\left(A_2(\frac{2}{3},s)\left|f'(a)\right|^q+A_3(\frac{2}{3},s)\left|f'(\frac{a+b}{2})\right|^q\right)^{\frac{1}{q}}\right. \\
&\left.+\left(A_2(\frac{2}{3},s)\left|f'(b)\right|^q+A_3(\frac{2}{3},s)\left|f'(\frac{a+b}{2})\right|^q\right)^{\frac{1}{q}}\right\},
\end{aligned}$$

where

$$\begin{aligned}
A_2(\frac{2}{3},s) &= \frac{2^{s+3}+3^{s+1}(s-1)}{3^{s+2}(s+1)(s+2)}, \\
A_3(\frac{2}{3},s) &= \frac{2+3^{s+1}(2s+1)}{3^{s+2}(s+1)(s+2)}.
\end{aligned}$$

Corollary 7. *Under the assumptions of Theorem 2 with* $\lambda = \frac{1}{2}$ *and* $\theta = 1$*, then we have following midpoint type inequality*

$$\begin{aligned}
&\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(x)dx\right| \leq \frac{b-a}{8}\left(\frac{2}{(s+1)(s+2)}\right)^{\frac{1}{q}} \\
&\times\left\{\left(\left|f'(a)\right|^q+(s+1)\left|f'(\frac{a+b}{2})\right|^q\right)^{\frac{1}{q}}\right. \\
&\left.+\left(\left|f'(b)\right|^q+(s+1)\left|f'(\frac{a+b}{2})\right|^q\right)^{\frac{1}{q}}\right\}.
\end{aligned}$$

Corollary 8. *Under the assumptions of Theorem 2 with* $\lambda = \frac{1}{2}$ *, and* $\theta = 0$*, then we get the following trapezoid type inequality*

$$\begin{aligned}
&\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)dx\right| \leq \frac{b-a}{8}\left(\frac{2}{(s+1)(s+2)}\right)^{\frac{1}{q}} \\
&\times\left\{\left((s+1)\left|f'(a)\right|^q+\left|f'(\frac{a+b}{2})\right|^q\right)^{\frac{1}{q}}\right. \\
&\left.+\left((s+1)\left|f'(b)\right|^q+\left|f'(\frac{a+b}{2})\right|^q\right)^{\frac{1}{q}}\right\}.
\end{aligned}$$

Using Lemma 1 we shall give another result for $s-$convex functions as follows.

Theorem 3. *Let* $f : I \subseteq [0,\infty) \rightarrow \mathbb{R}$ *be a differentiable mapping on* I° *such that* $f' \in L[a,b]$*, where* $a,b \in I^\circ$ *with* $a < b$ *and* $\theta,\lambda \in [0,1]$*. If* $|f'|^q$ *is* $s-$*convex on* $[a,b]$*,* $q > 1$*, then the following inequality holds:*

$$\left|(1-\theta)\left(\lambda f(a)+(1-\lambda)f(b)\right)+\theta f((1-\lambda)a+\lambda b)-\frac{1}{b-a}\int_a^b f(x)dx\right|$$

$$
\begin{aligned}
&\leq (b-a)\left(\frac{\theta^{p+1}+(1-\theta)^{p+1}}{p+1}\right)^{\frac{1}{p}} \\
&\quad \times\left[\lambda^{2}\left(\frac{\left|f'(a)\right|^{q}+\left|f'(C)\right|^{q}}{s+1}\right)^{\frac{1}{q}}+(1-\lambda)^{2}\left(\frac{\left|f'(b)\right|^{q}+\left|f'(C)\right|^{q}}{s+1}\right)^{\frac{1}{q}}\right]. \qquad (2.9)
\end{aligned}
$$

where $C=(1-\lambda)a+\lambda b$ *and* $\frac{1}{p}+\frac{1}{q}=1$.

Proof. Suppose that $C=(1-\lambda)a+\lambda b$. From Lemma 1 and by Hölder's integral inequality, we have

$$
\begin{aligned}
&\left|(1-\theta)(\lambda f(a)+(1-\lambda)f(b))+\theta f(C)-\frac{1}{b-a}\int_{a}^{b}f(x)dx\right| \leq (b-a) \\
&\left[\lambda^{2}\int_{0}^{1}|t-\theta|\left|f'(ta+(1-t)C)\right|dt+(1-\lambda)^{2}\int_{0}^{1}|t-\theta|\left|f'(tb+(1-t)C)\right|dt\right] \\
&\leq (b-a)\left\{\lambda^{2}\left(\int_{0}^{1}|t-\theta|^{p}dt\right)^{\frac{1}{p}}\left(\int_{0}^{1}\left|f'(ta+(1-t)C)\right|^{q}dt\right)^{\frac{1}{q}}\right. \\
&\quad \left.+(1-\lambda)^{2}\left(\int_{0}^{1}|t-\theta|^{p}dt\right)^{\frac{1}{p}}\left(\int_{0}^{1}\left|f'(tb+(1-t)C)\right|^{q}dt\right)^{\frac{1}{q}}\right\}. \qquad (2.10)
\end{aligned}
$$

Since $|f'|^{q}$ is $s-$convex on $[a,b]$, the inequalities (2.3) and (2.4) holds. Hence, by simple computation

$$
\begin{aligned}
\int_{0}^{1}\left|f'(ta+(1-t)C)\right|^{q}dt &\leq \int_{0}^{1}t^{s}\left|f'(a)\right|^{q}+(1-t)^{s}\left|f'(C)\right|^{q} \\
&= \frac{\left|f'(a)\right|^{q}+\left|f'(C)\right|^{q}}{s+1} \qquad (2.11)
\end{aligned}
$$

$$
\begin{aligned}
\int_{0}^{1}\left|f'(tb+(1-t)C)\right|^{q}dt &\leq \int_{0}^{1}t^{s}\left|f'(b)\right|^{q}+(1-t)^{s}\left|f'(C)\right|^{q} \\
&= \frac{\left|f'(b)\right|^{q}+\left|f'(C)\right|^{q}}{s+1} \qquad (2.12)
\end{aligned}
$$

and

$$
\int_{0}^{1}|t-\theta|^{p}dt=\frac{\theta^{p+1}+(1-\theta)^{p+1}}{p+1} \qquad (2.13)
$$

thus, using (2.11)-(2.13) in (2.10), we obtain the inequality (2.9). This completes the proof. □

Corollary 9. *Under the assumptions of Theorem 3 with $s=1$, we have*

$$\left|(1-\theta)\left(\lambda f(a)+(1-\lambda)f(b)\right)+\theta f((1-\lambda)a+\lambda b)-\frac{1}{b-a}\int\limits_a^b f(x)dx\right|$$

$$\begin{aligned}\leq\quad & (b-a)\left(\frac{\theta^{p+1}+(1-\theta)^{p+1}}{p+1}\right)^{\frac{1}{p}}\left(\frac{1}{2}\right)^{\frac{1}{q}}\\ & \times\left[\lambda^2\left(\left|f'(a)\right|^q+\left|f'(C)\right|^q\right)^{\frac{1}{q}}+(1-\lambda)^2\left(\left|f'(b)\right|^q+\left|f'(C)\right|^q\right)^{\frac{1}{q}}\right],\end{aligned}$$

where $C=(1-\lambda)a+\lambda b$ *and* $\frac{1}{p}+\frac{1}{q}=1$.

Corollary 10. *Under the assumptions of Theorem 3 with $\theta=1$, then we have the following generalized midpoint type inequality*

$$\left|f((1-\lambda)a+\lambda b)-\frac{1}{b-a}\int\limits_a^b f(x)dx\right|\leq(b-a)\left(\frac{1}{p+1}\right)^{\frac{1}{p}}$$

$$\times\left[\lambda^2\left(\frac{\left|f'(a)\right|^q+\left|f'(C)\right|^q}{s+1}\right)^{\frac{1}{q}}+(1-\lambda)^2\left(\frac{\left|f'(b)\right|^q+\left|f'(C)\right|^q}{s+1}\right)^{\frac{1}{q}}\right],$$

where $C=(1-\lambda)a+\lambda b$ *and* $\frac{1}{p}+\frac{1}{q}=1$.

Corollary 11. *Under the assumptions of Theorem 3 with $\theta=0$, then we have the following generalized trapezoid type inequality*

$$\left|\lambda f(a)+(1-\lambda)f(b)-\frac{1}{b-a}\int\limits_a^b f(x)dx\right|\leq(b-a)\left(\frac{1}{p+1}\right)^{\frac{1}{p}}$$

$$\times\left[\lambda^2\left(\frac{\left|f'(a)\right|^q+\left|f'(C)\right|^q}{s+1}\right)^{\frac{1}{q}}+(1-\lambda)^2\left(\frac{\left|f'(b)\right|^q+\left|f'(C)\right|^q}{s+1}\right)^{\frac{1}{q}}\right],$$

where $C=(1-\lambda)a+\lambda b$ *and* $\frac{1}{p}+\frac{1}{q}=1$.

Corollary 12. *Under the assumptions of Theorem 3 with $\theta=1$, if $|f'(x)|\leq M$, $x\in[a,b]$, then we have the following Ostrowski type inequality*

$$\left|f(x)-\frac{1}{b-a}\int\limits_a^b f(u)du\right|\leq\frac{M}{(p+1)^{\frac{1}{p}}}\left(\frac{2}{s+1}\right)^{\frac{1}{q}}\left[\frac{(x-a)^2+(b-x)^2}{b-a}\right]\tag{2.14}$$

for each $x\in[a,b]$.

Proof. For each $x\in[a,b]$, there exist $\lambda_x\in[0,1]$ such that $x=(1-\lambda_x)a+\lambda_x b$. Hence we have $\lambda_x=\frac{x-a}{b-a}$ and $1-\lambda_x=\frac{b-x}{b-a}$. Therefore for each $x\in[a,b]$, from the inequality (2.9) we obtain the inequality (2.14). □

Remark 2. *We note that the inequality (2.14) is the same of the inequality in* [2, Theorem 3].

Corollary 13. *Under the assumptions of Theorem 3 with* $\lambda = \frac{1}{2}$ *and* $\theta = \frac{2}{3}$, *then we have the following Simpson type inequality*

$$\left|\frac{1}{6}\left[f(a)+4f\left(\frac{a+b}{2}\right)+f(b)\right]-\frac{1}{b-a}\int\limits_a^b f(x)dx\right|$$

$$\leq \frac{b-a}{12}\left(\frac{1+2^{p+1}}{3(p+1)}\right)^{\frac{1}{p}}\left\{\left(\frac{\left|f'\left(\frac{a+b}{2}\right)\right|^q+\left|f'(a)\right|^q}{s+1}\right)^{\frac{1}{q}}+\left(\frac{\left|f'\left(\frac{a+b}{2}\right)\right|^q+\left|f'(b)\right|^q}{s+1}\right)^{\frac{1}{q}}\right\},$$

which is the same of the inequality in [12, Theorem 8].

Corollary 14. *Under the assumptions of Theorem 3 with* $\lambda = \frac{1}{2}$ *and* $\theta = 1$, *then we have the following midpoint type inequality*

$$\begin{aligned}\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int\limits_a^b f(x)dx\right| &\leq \frac{b-a}{4}\left(\frac{1}{p+1}\right)^{\frac{1}{p}} \\ &\times\left\{\left(\frac{\left|f'\left(\frac{a+b}{2}\right)\right|^q+\left|f'(a)\right|^q}{s+1}\right)^{\frac{1}{q}}+\left(\frac{\left|f'\left(\frac{a+b}{2}\right)\right|^q+\left|f'(b)\right|^q}{s+1}\right)^{\frac{1}{q}}\right\}.\end{aligned} \tag{2.15}$$

Remark 3. *We note that the inequality (2.15) is better than the inequality in* [3, Theorem 2.3]. *Because, by inequality*

$$2^{s-1}\left|f'\left(\frac{a+b}{2}\right)\right|^q \leq \frac{\left|f'(a)\right|^q+\left|f'(b)\right|^q}{s+1}$$

we have

$$\begin{aligned}\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int\limits_a^b f(x)dx\right| &\leq \left(\frac{b-a}{4}\right)\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left(\frac{1}{s+1}\right)^{\frac{2}{q}} \\ &\quad\times\left[\left(\left(2^{1-s}+s+1\right)\left|f'(a)\right|^q+2^{1-s}\left|f'(b)\right|^q\right)^{\frac{1}{q}}\right. \\ &\quad\left.+\left(\left(2^{1-s}+s+1\right)\left|f'(b)\right|^q+2^{1-s}\left|f'(a)\right|^q\right)^{\frac{1}{q}}\right],\end{aligned}$$

which is the same of the inequality in [3, Theorem 2.3].

Corollary 15. *Under the assumptions of Theorem 3 with* $\lambda = \frac{1}{2}$ *and* $\theta = 0$, *then we have the following trapezoid type inequality*

$$\begin{aligned}\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int\limits_a^b f(x)dx\right| &\leq \frac{b-a}{4}\left(\frac{1}{p+1}\right)^{\frac{1}{p}} \\ &\times\left\{\left(\frac{\left|f'\left(\frac{a+b}{2}\right)\right|^q+\left|f'(a)\right|^q}{s+1}\right)^{\frac{1}{q}}+\left(\frac{\left|f'\left(\frac{a+b}{2}\right)\right|^q+\left|f'(b)\right|^q}{s+1}\right)^{\frac{1}{q}}\right\}.\end{aligned} \tag{2.16}$$

We note that the obtained inequality (2.16) is better than the first inequality in [9, Theorem 3].

Theorem 4. *Let $f : I \subseteq [0,\infty) \to \mathbb{R}$ be a differentiable mapping on I° such that $f' \in L[a,b]$, where $a, b \in I^{\circ}$ with $a < b$ and $\alpha, \lambda \in [0,1]$. If $|f'|^{q}$ is $s-$concave on $[a,b]$, $q > 1$, then the following inequality holds:*

$$\left|(1-\theta)\left(\lambda f(a)+(1-\lambda)f(b)\right)+\theta f((1-\lambda)a+\lambda b)-\frac{1}{b-a}\int\limits_{a}^{b}f(x)dx\right| \tag{2.17}$$

$$\leq (b-a)\left(\frac{1}{2}\right)^{\frac{1-s}{q}}\left(\frac{\theta^{p+1}+(1-\theta)^{p+1}}{p+1}\right)^{\frac{1}{p}}$$

$$\times\left\{\lambda^{2}\left|f'\left(\frac{(2-\lambda)a+\lambda b}{2}\right)\right|+(1-\lambda)^{2}\left|f'\left(\frac{(1-\lambda)a+(1+\lambda)b}{2}\right)\right|\right\},$$

where $C = (1-\lambda)a+\lambda b$ and $1/p+1/q = 1$.

Proof. Suppose that $C = (1-\lambda)a+\lambda b$. We proceed similarly as in the proof Theorem 3. Since $|f'|^{q}$ is $s-$concave on $[a,b]$, $\lambda \in (0,1]$ by the inequality (1.3), we get

$$\frac{1}{C-a}\int\limits_{a}^{C}\left|f'(x)\right|^{q}dx$$

$$= \int\limits_{0}^{1}\left|f'(ta+(1-t)C)\right|^{q}dt \leq \frac{1}{2^{1-s}}\left|f'\left(\frac{(2-\lambda)a+\lambda b}{2}\right)\right|^{q} \tag{2.18}$$

the inequality (2.18) also holds $\lambda = 0$ too. Similarly, for $\lambda \in [0,1)$ by the inequality (1.3), we have

$$\frac{1}{b-C}\int\limits_{C}^{b}\left|f'(x)\right|^{q}dx$$

$$= \int\limits_{0}^{1}\left|f'(tb+(1-t)C)\right|^{q}dt \leq \frac{1}{2^{1-s}}\left|f'\left(\frac{(1-\lambda)a+(1+\lambda)b}{2}\right)\right|^{q} \tag{2.19}$$

the inequality (2.19) also holds $\lambda = 1$ too. Thus using (2.13), (2.18) and (2.19) in (2.10), we obtain the inequality (2.17). This completes the proof. □

Corollary 16. *Under the assumptions of Theorem 4 with $s = 1$, we have*

$$\left|(1-\theta)\left(\lambda f(a)+(1-\lambda)f(b)\right)+\theta f((1-\lambda)a+\lambda b)-\frac{1}{b-a}\int\limits_{a}^{b}f(x)dx\right|$$

$$\leq (b-a)\left(\frac{\theta^{p+1}+(1-\theta)^{p+1}}{p+1}\right)^{\frac{1}{p}}$$

$$\times\left\{\lambda^{2}\left|f'\left(\frac{(2-\lambda)a+\lambda b}{2}\right)\right|+(1-\lambda)^{2}\left|f'\left(\frac{(1-\lambda)a+(1+\lambda)b}{2}\right)\right|\right\}.$$

Remark 4. *In Corollary 16, if we take* $\lambda=\frac{1}{2}$ *and* $\theta=0$, *then we have the following trapezoid type inequality.*

$$\begin{aligned}&\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)dx\right|\\ \leq&\ \frac{b-a}{4}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left[\left|f'\left(\frac{3b+a}{4}\right)\right|+\left|f'\left(\frac{3a+b}{4}\right)\right|\right]\end{aligned}$$

which is the same of the inequality in [9, Theorem 2]

Remark 5. *In Corollary 16, if we take* $\lambda=\frac{1}{2}$ *and* $\theta=1$, *then we have the following midpoint type inequality*

$$\begin{aligned}&\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(x)dx\right|\\ \leq&\ \frac{b-a}{4}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left[\left|f'\left(\frac{3b+a}{4}\right)\right|+\left|f'\left(\frac{3a+b}{4}\right)\right|\right]\end{aligned}$$

which is the same of the inequality in [3, Theorem 2.5].

Corollary 17. *Under the assumptions of Theorem 4 with* $\theta=0$, *we have the following generalized trapezoid type inequality*

$$\begin{aligned}&\left|\lambda f(a)+(1-\lambda)f(b)-\frac{1}{b-a}\int_a^b f(x)dx\right|\\ \leq&\ (b-a)\left(\frac{1}{2}\right)^{\frac{1-s}{q}}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\\ &\times\left\{\lambda^2\left|f'\left(\frac{(2-\lambda)a+\lambda b}{2}\right)\right|+(1-\lambda)^2\left|f'\left(\frac{(1-\lambda)a+(1+\lambda)b}{2}\right)\right|\right\},\end{aligned}$$

Corollary 18. *Under the assumptions of Theorem 4 with* $\theta=1$, *we have the following generalized midpoint type inequality*

$$\begin{aligned}&\left|f((1-\lambda)a+\lambda b)-\frac{1}{b-a}\int_a^b f(x)dx\right|\\ \leq&\ (b-a)\left(\frac{1}{2}\right)^{\frac{1-s}{q}}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\\ &\times\left\{\lambda^2\left|f'\left(\frac{(2-\lambda)a+\lambda b}{2}\right)\right|+(1-\lambda)^2\left|f'\left(\frac{(1-\lambda)a+(1+\lambda)b}{2}\right)\right|\right\},\end{aligned}$$

Corollary 19. *Under the assumptions of Theorem 3 with* $\theta=1$, *we have the following Ostrowski type inequality*

$$\begin{aligned}&\left|f(x)-\frac{1}{b-a}\int_a^b f(u)du\right|\\ \leq&\ \frac{2^{\frac{s-1}{q}}}{(p+1)^{\frac{1}{p}}(b-a)}\left[(x-a)^2\left|f'\left(\frac{x+a}{2}\right)\right|+(b-x)^2\left|f'\left(\frac{x+b}{2}\right)\right|\right]\end{aligned}\tag{2.20}$$

for each $x \in [a,b]$ *.*

Proof. For each $x \in [a,b]$, there exist $\lambda_x \in [0,1]$ such that $x = (1-\lambda_x)\, a + \lambda_x b$. Hence we have $\lambda_x = \frac{x-a}{b-a}$ and $1-\lambda_x = \frac{b-x}{b-a}$. Therefore for each $x \in [a,b]$, from the inequality (2.17) we obtain the inequality (2.20). □

Remark 6. *We note that the inequality (2.20) is the same of the inequality in* [2, Theorem 5]*.*

Corollary 20. *Under the assumptions of Theorem 4 with* $\lambda = \frac{1}{2}$ *and* $\theta = 0$*, we have the following trapezoid type inequality*

$$\left|\frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)dx\right| \le \frac{b-a}{4}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left(\frac{1}{2}\right)^{\frac{1-s}{q}}\left[\left|f'\left(\frac{3b+a}{4}\right)\right| + \left|f'\left(\frac{3a+b}{4}\right)\right|\right] \tag{2.21}$$

which is the same of the inequality in [10, Theorem 8 (i)]*.*

Corollary 21. *Under the assumptions of Theorem 4 with* $\lambda = \frac{1}{2}$ *and* $\theta = 1$*, we have the following midpoint type inequality*

$$\left|f\left(\frac{a+b}{2}\right) - \frac{1}{b-a}\int_a^b f(x)dx\right| \le \frac{b-a}{4}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left(\frac{1}{2}\right)^{\frac{1-s}{q}}\left[\left|f'\left(\frac{3b+a}{4}\right)\right| + \left|f'\left(\frac{3a+b}{4}\right)\right|\right] \tag{2.22}$$

which is the same of the inequality in [10, Theorem 8 (ii)]*.*

Corollary 22. *Under the assumptions of Theorem 4 with* $\lambda = \frac{1}{2}$ *and* $\theta = \frac{2}{3}$*, we have the following trapezoid type inequality*

$$\left|\frac{1}{6}\left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b)\right] - \frac{1}{b-a}\int_a^b f(x)dx\right| \le \frac{b-a}{12}\left(\frac{1+2^{p+1}}{3(p+1)}\right)^{\frac{1}{p}} 2^{\frac{s-1}{q}}\left[\left|f'\left(\frac{3b+a}{4}\right)\right| + \left|f'\left(\frac{3a+b}{4}\right)\right|\right]. \tag{2.23}$$

Remark 7. *If* $|f'|^q$ *,* $q>1$*, is concave on* $[a,b]$ *, using the power mean inequality, we have*

$$\begin{aligned} |f'(\lambda x + (1-\lambda) y)|^q &\ge \lambda |f'(x)|^q + (1-\lambda)|f'(y)|^q \\ &\ge (\lambda |f'(x)| + (1-\lambda)|f'(y)|)^q , \end{aligned}$$

$\forall x, y \in [a,b]$ *and* $\lambda \in [0,1]$ *. Hence*

$$|f'(\lambda x + (1-\lambda) y)| \ge \lambda |f'(x)| + (1-\lambda)|f'(y)|$$

so $|f'|$ *is also concave. Then by the inequality (1.1), we have*

$$\left|f'\left(\frac{3b+a}{4}\right)\right| + \left|f'\left(\frac{3a+b}{4}\right)\right| \le 2\left|f'\left(\frac{a+b}{2}\right)\right|. \tag{2.24}$$

Thus, using the inequality (2.24) in (2.21), (2.22) and (2.23) we get

$$\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)dx\right| \leq \frac{b-a}{2}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left(\frac{1}{2}\right)^{\frac{1-s}{q}}\left|f'\left(\frac{a+b}{2}\right)\right|,$$

$$\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(x)dx\right| \leq \frac{b-a}{2}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left(\frac{1}{2}\right)^{\frac{1-s}{q}}\left|f'\left(\frac{a+b}{2}\right)\right|,$$

and

$$\left|\frac{1}{6}\left[f(a)+4f\left(\frac{a+b}{2}\right)+f(b)\right]-\frac{1}{b-a}\int_a^b f(x)dx\right|$$
$$\leq \frac{b-a}{12}\left(\frac{1+2^{p+1}}{3(p+1)}\right)^{\frac{1}{p}}\left(\frac{1}{2}\right)^{\frac{1-s}{q}}\left|f'\left(\frac{a+b}{2}\right)\right|,$$

respectively, where $s\in(0,1]$ *and* $1/p+1/q=1$.

3. Some applications for special means

in [8], the following example is given.

Let $s\in(0,1)$ and $a,b,c\in\mathbb{R}$.We define function $f:[0,\infty)\to\mathbb{R}$ as

$$f(t)=\begin{cases} a, & t=0\\ bt^s+c, & t>0\end{cases}.$$

If $b\geq 0$ and $0\leq c\leq a$, then $f\in K_s^2$. Hence, for $a=c=0$, $b=s+1$, $s\in\left(0,\frac{1}{q}\right)$, $q\geq 1$, we have $f:[0,\infty)\to[0,\infty)$, $f(t)=t^{s+1}$, $|f'|^q\in K_s^2$.

Let us recall the following special means of arbitrary real numbers a,b with $a\neq b$ and $\alpha\in[0,1]$:

(1) The weighted arithmetic mean

$$A_\alpha(a,b):=\alpha a+(1-\alpha)b,\ a,b\in\mathbb{R}.$$

(2) The unweighted arithmetic mean

$$A(a,b):=\frac{a+b}{2},\ a,b\in\mathbb{R}.$$

(3) The Logarithmic mean

$$L(a,b):=\frac{b-a}{\ln b-\ln a},\quad a\neq b,\quad a,b>0.$$

(4) Then $p-$Logarithmic mean

$$L_p(a,b):=\left(\frac{b^{p+1}-a^{p+1}}{(p+1)(b-a)}\right)^{\frac{1}{p}},\ p\in\mathbb{R}\backslash\{-1,0\},\ a,b>0.$$

Now, using the results of Section 2, some new inequalities are derived for the above means.

Proposition 1. *Let* $a, b \in \mathbb{R}$ *with* $0 < a < b,\ q \geq 1,\ s \in \left(0, \frac{1}{q}\right)$ *and* $\lambda, \theta \in [0,1]$ *we have the following inequality:*

$$\begin{aligned} & \left|(1-\theta)\, A_\lambda\left(a^{s+1}, b^{s+1}\right) + \theta A_\lambda^{s+1}(a,b) - L_{s+1}^{s+1}(a,b)\right| \\ \leq\ & (b-a)\, A_1^{1-\frac{1}{q}}(\theta)\,(s+1)\left\{\lambda^2\left[a^{sq} A_2(\theta,s) + A_\lambda^{sq}(b,a) A_3(\theta,s)\right]^{\frac{1}{q}}\right. \\ & \left. + (1-\lambda)^2\left[b^{sq} A_2(\theta,s) + A_\lambda^{sq}(b,a) A_3(\theta,s)\right]^{\frac{1}{q}}\right\} \end{aligned}$$

where $A_1(\theta),\ A_2(\theta,s),\ A_3(\theta,s)$ *are defined as in Theorem 2.*

Proof. The assertion follows from Theorem 2 for the function $f(t) = t^{s+1}$, $t \in [0,\infty)$, $s \in \left(0, \frac{1}{q}\right)$. ☐

Proposition 2. *Let* $a, b \in \mathbb{R}$ *with* $0 < a < b,\ p, q > 1,\ \frac{1}{p} + \frac{1}{q} = 1,\ s \in \left(0, \frac{1}{q}\right)$ *and* $\lambda, \theta \in [0,1]$ *we have the following inequality:*

$$\begin{aligned} & \left|(1-\theta)\, A_\lambda\left(a^{s+1}, b^{s+1}\right) + \theta A_\lambda^{s+1}(a,b) - L_{s+1}^{s+1}(a,b)\right| \\ \leq\ & (b-a)\left(\frac{\theta^{p+1} + (1-\theta)^{p+1}}{p+1}\right)^{\frac{1}{p}} (s+1)^{1-\frac{1}{q}} \\ & \times\left[\lambda^2\left(a^{sq} + A_\lambda^{sq}(b,a)\right)^{\frac{1}{q}} + (1-\lambda)^2\left(b^{sq} + A_\lambda^{sq}(b,a)\right)^{\frac{1}{q}}\right]. \end{aligned}$$

where $1/p + 1/q = 1$.

Proof. The assertion follows from Theorem 3 for the function $f(t) = t^{s+1}$, $t \in [0,\infty)$, $s \in \left(0, \frac{1}{q}\right)$. ☐

References

[1] Alomari, M., Darus, M., Dragomir, S.S. *New inequalities of Simpson's Type for $s-$convex functions with applications*, RGMIA Res. Rep. Coll. 12 (4) (2009) Article 9. Online http://ajmaa.org/RGMIA/v12n4.php.

[2] Alomaria, M., Darus, M., Dragomir, S.S., Cerone, P. *Ostrowski type inequalities for functions whose derivatives ares-convex in the second sense*, Applied Mathematics Letters 23 (2010) 1071–1076.

[3] Alomari, M.W., Darus, M., Kirmaci, U.S. *Some inequalities of Hermite-Hadamard type for $s-$convex functions*, Acta Math. Scientia, 31B (4) (2011)1643-1652.

[4] Breckner, W.W. *Stetigkeitsaussagen für eine Klasse verallgemeinerter konvexer funktionen in topologischen linearen Räumen*, Publ. Inst. Math. 23 (1978) 13–20.

[5] Dragomir, S.S., Fitzpatrik, S. *The Hadamard's inequality for $s-$convex functions in the second sense*, Demonstration Math.32(4), (1999), 687–696.

[6] Dragomir, S.S., Pearce, C.E.M. *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, Victoria University, 2000.

[7] Dragomir, S.S., Rassias, Th. M. *Ostrowski type inequalities and applications in numerical integration*, Kluwer Academic Publishers, Dorcdrecht, Boston, London, 2002.

[8] Hudzik, H., Maligranda, L. *Some remarks on $s-$convex functions*, Aequationes Math. 48 (1994) 100-111.

[9] Kirmaci, U.S., Bakula, M.K., Özdemir, M.E., Pečarić, J. *Hadamard-type inequalities for $s-$convex functions*, Applied Mathematics and Computation 193 (2007) 26-35.

[10] Park, J. *Hermite-Hadamard-type inequalities for real $\alpha-$star $s-$convex mappings*, J. Appl. Math. & Informatics, 28 (2010) No. 5-6, 1507-1518.

[11] Sarikaya, M.Z., Aktan, N. *On the generalization of some integral inequalities and their applications*, Mathematical and Computer Modelling, 54 (2011) 2175-2182.

[12] Sarikaya, M.Z., Set, E., Özdemir, M.E. *On new inequalities of Simpson's type for* $s-$*convex functions*, Computers and Mathematics with Applications 60 (2010) 2191-2199.

[13] Sarikaya, M.Z., Set, E., Özdemir, M.E. *On new inequalities of Simpson's type for convex functions*, RGMIA Res. Rep. Coll. 13 (2) (2010) Article 2.

DEPARTMENT OF MATHEMATICS, FACULTY OF ARTS AND SCIENCES,, GIRESUN UNIVERSITY, 28100, GIRESUN, TURKEY.

E-mail address: `imdat.iscan@giresun.edu.tr`